



# Education

KwaZulu-Natal Department of Education

**PHYSICAL SCIENCES P2 (CHEMISTRY)**

**PREPARATORY EXAMINATION**

**SEPTEMBER 2018**

**MEMORANDUM**

**NATIONAL  
SENIOR CERTIFICATE**

**GRADE 12**

**MARKS : 150**

**This memorandum consists of 9 pages.**

**The marking guidelines as per 2014 Examination Guidelines, pages 34-37 must be applied when marking this Paper.**

**QUESTION 1**

- 1.1 B ✓✓ (2)
- 1.2 C ✓✓ (2)
- 1.3 C ✓✓ (2)
- 1.4 A ✓✓ (2)
- 1.5 B ✓✓ (2)
- 1.6 B ✓✓ (2)
- 1.7 D ✓✓ (2)
- 1.8 A ✓✓ (2)
- 1.9 A ✓✓ (2)
- 1.10 C ✓✓ (2)
- [20]**

**QUESTION 2**

- 2.1.1 E ✓ (1)
- 2.1.2 B ✓ (1)
- 2.1.3 D ✓ (1)
- 2.1.4 F ✓ (1)
- 2.1.5 G ✓ (1)
- 2.2.1 2,4,4-trimethylpent-2-ene ✓✓ (2)
- 2.2.2  $C_nH_{2n}$  ✓ (1)
- 2.3.1 ethanol ✓ (1)
- 2.3.2 sulphuric acid ✓ (1)
- [10]**


**QUESTION 3**

- 3.1.1 a series of organic compounds that can be described by the same general formula ✓  
in which one member differs from the next with a CH<sub>2</sub> group. ✓ (2)
- 3.1.2 the temperature at which the vapour pressure equals atmospheric/external pressure. ✓✓ (2 or 0) (2)
- 3.2 C ✓  
As the boiling point increases the vapour pressure decreases. ✓  
C has the highest boiling point. ✓ (3)
- 3.3 B ✓ (1)
- 3.4.1 118,50 °C ✓ (1)
- 3.4.2 In addition to London forces and dipole-dipole forces, C has two sites for hydrogen bonding between the molecules ✓ resulting in the strongest intermolecular forces occurring between molecules of C. ✓  
The intermolecular forces between molecules of C require the most amount of energy to overcome. ✓  
C will therefore have the highest boiling point. ✓ (4)

**[13]****QUESTION 4**

- 4.1.1 Addition/hydrohalogenation ✓ (1)
- 4.1.2 Substitution/hydrolysis ✓ (1)

4.2




- Whole structure correct: 2/2
- Only functional group correct 1/2
- More than one functional group 0/2

2-bromobutane ✓

(3)

- 4.3 Secondary ✓  
The carbon to which the —O—H ✓ is bonded to, is bonded to TWO other carbon atoms. ✓ (3)
- 4.4 Dehydration ✓✓ (1)
- 4.5 (Gentle) heat ✓  
Aqueous/dilute strong base (accept NaOH(dilute) or KOH(dilute) ✓ (2)
- 4.6.1 Compounds with the same molecular formula, ✓ but different positions of the side chain/substituents/functional groups on the parent chain. ✓ (2)
- 4.6.2 Elimination ✓ (1)
- 4.6.3



1 mark for each reactant and product

(3)

**[17]**

**QUESTION 5****5.1.1 ANY ONE**

- The change in concentration ✓ of reactants/products per unit time. ✓
- Rate of ✓ change in concentration of reactants or products. ✓
- Change in amount/number of moles/volume/mass of reactants/products ✓ per (unit) time. ✓
- Amount/number of moles/volume/mass of products formed OR reactants used ✓ per (unit) time. ✓ (2)

5.1.2 60 - 61(s) ✓ (1)

5.1.3  $n(\text{CO}_2) = n(\text{CaCO}_3)$  ✓

$$= \left( n = \frac{m}{M} \right)$$

$$= \left( \frac{86 - 40}{100} \right) \checkmark$$

$$= 0,46 \text{ mols}$$

$$n = \frac{V}{V_m}$$

$$0,46 \checkmark = \frac{V}{22,4} \checkmark$$

$$V = 10,304 \text{ dm}^3 \checkmark \quad (5)$$

5.1.4 40 g ✓ (1)

5.1.5 INCREASES ✓ (1)

5.1.6 See attached graph. (3)

- Curve starts at 86 g and ends at 40g ✓
- The completion time is above 60 or 61s ✓
- The curve above the original ✓

5.2.1 Collision theory ✓ (1)

5.2.2 The shaded areas in the distribution curves represent the number of molecules with sufficient kinetic energy to overcome the activation energy ✓. An increase in the temperature of the system results in a greater number of particles with sufficient kinetic energy to overcome the activation energy of the reaction ✓. This results in more effective collisions per unit time OR a higher chance of an effective collision occurring ✓, resulting in a higher reaction rate. (3)

**[17]****QUESTION 6**

6.1 When the equilibrium in a closed system is disturbed ✓, the system will re-instate a new equilibrium by favouring the reaction that will oppose the disturbance. ✓ (2)

6.2.1 REMAINS THE SAME ✓ (1)

6.2.2 INCREASES ✓ (1)

6.2.3 REMAINS THE SAME ✓ (1)

6.2.4 INCREASES ✓ (1)

**Apply negative marking from 6.2.4**

6.3 According to Le Chatelier's Principle a decrease in temperature favours the exothermic reaction ✓  
A decrease in temperature increases the equilibrium constant ✓. Therefore the forward reaction is favoured ✓ (3)

6.4

**Marking criteria:**

- Indicating that the number of mols of H<sub>2</sub> decreases by an unknown amount ✓
- Correct mol ratio ✓
- Calculating in terms of x the quantity(mol) at equilibrium of all three substances ✓
- Substitute  $V = 4 \text{ dm}^3$  in  $c = \frac{n}{V}$  to determine concentration at equilibrium of H<sub>2</sub>/I<sub>2</sub> and HI. ✓
- K<sub>c</sub> expression ✓
- Substitution of concentrations in K<sub>c</sub> expression ✓
- Substitution of 49 for K<sub>c</sub> ✓
- Equation:  $n = \frac{m}{M}$  ✓
- Substituting in the above equation ✓
- Final answer: 399,36 g ✓

No K<sub>c</sub> expression, correct substitution: Max.  $\frac{9}{10}$

Wrong K<sub>c</sub> expression : Max.  $\frac{6}{10}$

	H <sub>2</sub>	I <sub>2</sub>	HI	
Initial quantity(mol)	2	2	0	
Change(mol)	-x ✓	-x	+2x	Ratio ✓
Quantity at equilibrium(mol)	2-x	2-x	2x	✓
Equilibrium concentration(mol.dm <sup>-3</sup> )	$\frac{2-x}{4}$	$\frac{2-x}{4}$	$\frac{x}{2}$	Divide by 4 ✓

$$K_c = \frac{[HI]^2}{[H_2][I_2]} \checkmark = \frac{\left(\frac{x}{2}\right)^2}{\left(\frac{2-x}{4}\right)\left(\frac{2-x}{4}\right)} \checkmark = 49 \checkmark$$

$$\begin{aligned} x &= 1,56 \text{ mol} \\ m(\text{HI}) &= nM \checkmark \\ &= (2)(1,56)(128) \checkmark \\ &= 399,36 \text{ g} \checkmark \end{aligned}$$

(10)  
[19]

**QUESTION 7**

7.1 It dissociates/ionises completely in water to form a high concentration of OH<sup>-</sup> ions. ✓ (1)

7.2 It contains a small amount (number of moles) of base ✓ in proportion to the volume of water ✓ (2)

7.3

- Formula pH = -log [H<sub>3</sub>O<sup>+</sup>] ✓ / pOH = -log [OH<sup>-</sup>] ✓
- Substitute 13,45 for pH ✓ / 0,55 for pOH ✓
- c(OH<sup>-</sup>) = 0,282 mol.dm<sup>-3</sup> ✓
- Using ratio of 1: 2 to calculate c((Ba(OH)<sub>2</sub>) ✓
- Formula m = cVM ✓
- Substituting into the above formula ✓
- Answer ✓

**Option 1:**

$$\text{pH} = -\log [\text{H}_3\text{O}^+] \checkmark$$

$$13,45 \checkmark = -\log [\text{H}_3\text{O}^+]$$

$$\therefore [\text{H}_3\text{O}^+] = 3,54 \times 10^{-14} \text{ mol.dm}^{-3}$$

$$[\text{H}_3\text{O}^+][\text{OH}^-] = 1 \times 10^{-14}$$

$$c(\text{OH}^-) = 0,282 \text{ mol.dm}^{-3} \checkmark$$

$$c((\text{Ba}(\text{OH})_2) = 0,141 \text{ mol.dm}^{-3} \checkmark$$

$$m = cVM \checkmark$$

$$= \frac{(0,141)(0,25)(171)}{1} \checkmark$$

$$= 6,03 \text{ g} \checkmark$$

**Option 2:**

$$\text{pOH} = -\log [\text{OH}^-] \checkmark$$

$$0,55 \checkmark = -\log [\text{OH}^-]$$

$$\therefore [\text{OH}^-] = 0,282 \text{ mol.dm}^{-3} \checkmark$$

$$c((\text{Ba}(\text{OH})_2) = 0,141 \text{ mol.dm}^{-3} \checkmark$$

$$m = cVM \checkmark$$

$$= \frac{(0,141)(0,25)(171)}{1} \checkmark$$

$$= 6,03 \text{ g} \checkmark$$

(7)

7.4 Positive marking from question 7.3: concentration of Ba(OH)<sub>2</sub>

**Marking guidelines**

- Formulae:  $c = \frac{n}{V} / n = cV / \frac{c_a \times V_a}{c_b \times V_b} = \frac{n_a}{n_b} \checkmark$
- Substitution of:  $0,141 \times 60 / 0,141 \times 0,06 \checkmark$
- Use mol ratio:  $n_a : n_b = 2 : 1 \checkmark$
- Final answer:  $33,84 \text{ cm}^3 / 0,03384 \text{ dm}^3 \checkmark$

**Option 1:**

$$n(\text{HCl}) = 2n((\text{Ba}(\text{OH})_2)$$

$$= 2cV$$

$$= \frac{2(0,141)(0,06)}{1} \checkmark$$

$$= 0,01692 \text{ mols}$$

$$c(\text{HCl}) = n/V \checkmark$$

$$0,5 \checkmark = 0,01692/V$$

$$V = 0,03384 \text{ dm}^3 / 33,84 \text{ cm}^3 \checkmark$$

**Option 2:**

$$\frac{c_A}{c_B} \frac{V_A}{V_B} = \frac{n_A}{n_B} \checkmark$$

$$\frac{0,5}{0,141} \frac{V_A}{0,06} \checkmark = \frac{2}{1} \checkmark$$

$$V_A = 0,03384 \text{ dm}^3 \checkmark$$

$$\text{Accept } V_B = 60 \text{ cm}^3$$

$$V_A = 33,84 \text{ cm}^3$$

(4)

[14]

**QUESTION 8**

8.1 GALVANIC, ✓ converts chemical energy to electrical energy ✓ or no dc power supply. (2)

8.2 Temperature of 25 °C/298K ✓  
Pressure 101,3 kPa ✓Concentration of electrolyte of 1 mol.dm<sup>-3</sup> ✓ (3)


8.3 Chlorine (molecule) ✓ ✓ (2)

8.4 **OPTION 1**

$$E^{\ominus}_{\text{cell}} = E^{\ominus}_{\text{cathode}} - E^{\ominus}_{\text{anode}} \checkmark$$

$$= 1,36 \checkmark - (-1,66) \checkmark$$

$$= 3,02 \text{ V} \checkmark$$

**OPTION 2****Notes**

- Accept any other correct formula from the data sheet.
- Any other formula using unconventional abbreviations, e.g.  $E^{\ominus}_{\text{cell}} = E^{\ominus}_{\text{OA}} - E^{\ominus}_{\text{RA}}$  followed by correct substitutions:

$$E^{\ominus}_{\text{sel}} = E^{\ominus}_{\text{OM}} - E^{\ominus}_{\text{RM}} \text{ Max: } \frac{3}{4}$$

8.5  $3\text{Cl}_2(\text{g}) + 2\text{Al}(\text{s}) \rightarrow 6\text{Cl}^-(\text{aq}) + 2\text{Al}^{3+}(\text{aq})$ **Notes**

- Reactants ✓          Products ✓          Balancing ✓
- Ignore phases.
- Marking rule 6.3.10
- Marking rule 3.9.
- Marking rule 3.4: One mark is forfeited when the charge of an ion is omitted per equation (not for the charge on the electron)

(3)

8.6.1 REMAINS THE SAME ✓

(1)

8.6.2 DECREASES ✓

(1)

**[16]****QUESTION 9**

9.1 A solution that conducts electricity through the movement of ions. ✓ (1)

9.2  $\text{Cu}^{2+}$  ✓ (1)

9.3.1 Decreases ✓ (1)

9.3.2  $\text{Cu} \rightarrow \text{Cu}^{2+} + 2\text{e}^-$ **Notes**

- $\text{Cu}^{2+} + 2\text{e}^- \leftarrow \text{Cu} \quad (\frac{2}{2})$            $\text{Cu}^{2+} + 2\text{e}^- \rightleftharpoons \text{Cu} \quad (\frac{0}{2})$
- $\text{Cu} \rightleftharpoons \text{Cu}^{2+} + 2\text{e}^- \quad (\frac{1}{2})$            $\text{Cu}^{2+} + 2\text{e}^- \rightarrow \text{Cu} \quad (\frac{0}{2})$
- Ignore if charge on electron is omitted.
- If a charge of an ion is omitted e.g.  $\text{Cu} \rightarrow \text{Cu}^2 + 2\text{e}^-$  is  $\text{Cu} \rightarrow \text{Cu}^2 + 2\text{e}^-$  Max.:  $\frac{1}{2}$

(2)

9.3.4

**Marking criteria**

- Calculate number of mols of cations:  $2,259 \times 10^{24} = n(6,023 \times 10^{23})$  ✓
- Formula:  $n = \frac{m}{M}$  ✓
- Substitute calculated number of moles of cations and 63,5 in  $n = \frac{m}{M}$  ✓
- Final answer 238,125 g ✓

$$n_e = nNA$$

$$2,259 \times 10^{24} = n(6,023 \times 10^{23}) \quad \checkmark$$

$$n = 3,75 \text{ mols}$$

$$m = nM \quad \checkmark$$

$$= (3,75)(63,5) \quad \checkmark$$

$$= 238,125 \text{ g.} \quad \checkmark$$

(4)  
[9]**QUESTION 10**

- 10.1.1 Haber ✓ (1)
- 10.1.2 Catalytic oxidation of ammonia ✓ (1)
- 10.1.3 Nitrogen dioxide ✓ (1)
- 10.1.4 Ammonium nitrate ✓ (1)
- 10.2.1 Sulphuric acid/H<sub>2</sub>SO<sub>4</sub> ✓ (1)
- 10.2.2  $\text{H}_2\text{SO}_4 + 2\text{NH}_3 \rightarrow (\text{NH}_4)_2\text{SO}_4$

**Notes:**

- Reactants ✓ Products ✓ Balancing ✓
- Marking rule 6.3.10.

(3)

$$\begin{aligned} 10.3 \quad \% \text{ N} &= 14/20 \times 36 \\ &= 25,2\% \quad \checkmark \end{aligned}$$

$$\text{Mass of N} = 25,2/100 \times m$$

$$12,60 \checkmark = \frac{25,2}{100} \times m \quad \checkmark$$

$$m = 50 \text{ kg} \quad \checkmark$$

(4)

10.4 Fertiliser A ✓

Fertilizer A has a high percentage of Phosphorus compared to fertilizer B. ✓✓

(3)

[15]

**TOTAL MARKS: 150**



**QUESTION 5.1.6**